



## Knowledge, Awareness and Preventive Practices of Dengue in Pakistan

Atija Waris<sup>1</sup>, Muhammad Mufeez Sajjad<sup>2</sup>, Talal Ehsan<sup>3</sup>, Sana Hafeez<sup>4</sup>, Umar Sultan<sup>5</sup>,  
Naglaa M Shalaby<sup>6</sup>

<sup>1</sup>Institute of Molecular Biology and Biotechnology, Zoology, University of Lahore, Punjab, Pakistan.

<sup>2</sup>Government Graduate College Ravi Road, Lahore, Punjab, Pakistan

<sup>3</sup>Department of Medicine, Shifa Hospital Islamabad, Pakistan.

<sup>4</sup>University of Management and Technology, Lahore, Punjab, Pakistan.

<sup>5</sup>Department of Cardiology, Punjab Institute of Cardiology Lahore, Pakistan.

<sup>6</sup>Department of Microbiology, Faculty of Medicine, Northern Border University, Arar 91431, Saudi Arabia.

### ARTICLE INFO

#### Keywords

Dengue, Fever, School, Education, Awareness, Knowledge

**Corresponding Author:** Sana Hafeez,  
University of Management and Technology,  
Lahore, Punjab, Pakistan.  
Email: [Drsana140@gmail.com](mailto:Drsana140@gmail.com)

#### Declaration

**Authors' Contribution:** All authors equally contributed to the study and approved the final manuscript.

**Conflict of Interest:** No conflict of interest.

**Funding:** No funding received by the authors.

#### Article History

Received: 13-12-2024, Revised: 02-02-2025

Accepted: 08-03-2025, Published: 31-03-2025

### ABSTRACT

**Objective:** Dengue, a mosquito-borne disease has become endemic. This study was conducted to explore the knowledge, awareness and preventive practices of dengue in the Lahore district. This study was quantitative and a cross-sectional survey was conducted in the study area. **Material and methods:** A total of 160 randomly chosen respondents from the 20 public secondary schools participated in the survey. Data were collected through a structured, validated and reliable questionnaire. Collected data were analyzed using the Statistical Package of Social Sciences (SPSS). Descriptive statistics were applied to the data to explore the distribution of responses regarding knowledge, awareness and prevention practices against dengue. **Results:** The majority (86.9%) of respondents had heard about dengue fever and for 90.6% of respondents, Aedes was the carrier of dengue. Around 56% of respondents had believed that Aedes breed in unclean water and 96.2% had a mistaken belief that it bites often at noon. Fever was the leading symptom (80.6%) and the use of mosquito repellent coil (98.8%), net (95%) and lotion (88.1%) were key preventive measures. Almost 76% of respondents were informed to visit the nearest Basic Health Unit (BHU) for urgent treatment followed by 21.2% of respondents favoring consulting a Hakeem (quacks) for the treatment. **Conclusion:** The school was the leading information source (80.6%) and the role of modern gadgets in awareness creating was dismal. This is suggested that community awareness level is obligatory, integration of modern gadgets in awareness campaigns is requisite, thus for high preventive measures synergistic strategy is needed.

### INTRODUCTION

Dengue is one of the prominent threats to society tending to cause a large number of human threats. Dengue, the mosquito-borne disease, transmitted by the bites of Aedes mosquitoes, primarily Aedes aegypti and Aedes albopictus, is considered the most widespread human arboviral infection across the world (Sharmin et al., 2015). Brady et al. (2012) have reported 20,000 deaths each year due to dengue in the world. The cause of dengue fever (DF) is the infection with any one of the 4 serotypes (DENV-1, 2, 3, and 4) of dengue virus and the DF may appear as a fatal disease characterized by dengue hemorrhagic fever (DHF) and dengue shock syndrome (DSS) (Lambrechts et al., 2010; Naing et al., 2011). Dengue fever is caused by a virus named single standard RNA by biting three types of genus Aedes and Alboictus (Hafeez et al., 2012). Dengue fever (DF) is responsible for significant human diseases and deaths

worldwide (Bhatt, 2013). Aedes aegypti is known as the primary vector for DF in both rural and urban areas of the world (Lambrechts et al., 2010). Many species of mosquitoes affect people and causes many diseases in human such as malaria, yellow fever and dengue fever also (Murugan et al., 2007).

The first dengue virus case emerged in South East Asia and approximately 52% of people were vulnerable to the risk of dengue around the world (Vachvanichsanong and Thisyakorn, 2016). Pakistan lies in the south Asian region and is also considered one of the prominent countries for its diverse geographical circumference. The first confirmed case of dengue fever was reported in 1994 in Pakistan. However, the annual prevalent trend and startling rise in dengue cases first became ostensible in November 2005 (WHO, 2021; Jehan, 2011). Since 2010, Pakistan has seen an

incredible number of cases of dengue. In 2020, 47,120 confirmed cases were reported of which 75 casualties were recorded as well. By the end of 2021, 48,906 dengue cases including 183 deaths were recorded (Dengue fever, 2022). Over the period and pertinent to its hefty penetration the dengue has become endemic in Pakistan mingling the whole year with a high level of prevalence in the post moon soon season expanding the vulnerability across the four provinces (Jahan, 2011).

The health sector of Pakistan has reported the lethal effects and fatalities of dengue in Pakistan for reasons including unhealthy food, and edibles with unsuitable sanitation in the community (Ali et al., 2017). According to Islam et al. (2022), owing to deprived healthcare capacity, slower vaccination rate, increasing COVID-19 variants, socio-economic inequalities, and misinformation the dengue fatalities are mounting over time.

DF is an automatic fever, which usually lasts for 5-7 days. It is sometimes weakened during the acute phase of the disease. The medical characteristics of DF vary according to the age of the patient. Newborns and young children may have an uncontrolled feral disease with mycoplasma rash. Older children and adults may have mild fever syndrome or high fever (usually biphasic), retroviral pain, myalgia, arthralgia, nausea, vomiting, and petechiae. Leukopenia and thrombocytopenia are common in all ages. In some cases, DF bleeding may be accompanied by complications such as gangrene, epilepsy, gastrointestinal bleeding, and hematuria (Swaminathan et al., 2013).

Numerous factors such as high population growth, rapid urbanization, worsening of the waste management system (Siddiqua et al., 2018) and lack of vector control accelerated the expansion and transmission of dengue (Jeelani et al., 2015). Inadequate water supply (Jeelani et al., 2015), illiteracy, poverty and social disparities have an association with poor dengue management (Guzma'n and Kouri, 2002).

Currently, no effective vaccine against dengue is available, this the only possible method of prevention of dengue is vector control (Udayanga et al., 2018). However, for this prevention method community participation and awareness is obligatory (Naing et al., 2011). Effective participation of the community heavily depends on knowledge, awareness and attitude of people toward the particular disease (Jeelani et al., 2015). In this preview, effective dengue prevention and remedy dengue has become a key concern in Pakistan. Therefore, this study was conducted with the objective aiming at assessing the awareness knowledge and attitude of students towards dengue prevention.

## METHODOLOGY

This study was conducted in rural areas of District Lahore. Lahore district is located between a longitude angle of 73 to 74 East and a latitude angle of 30 to 31.5 North with 600ft regular plains as elevation in north East of Punjab. The district covers 1,280 km<sup>2</sup> having six Tehsils named Lahore City, Lahore Sadar, Jaranwala, Chak Jhumrah, Samundari and Tandlianwala. Most of the urban area is located in Tehsil Lahore City, while the other five tehsils are consisting of rural areas (Bashir et al., 2010).

The survey method was used to collect data from the selected respondents. The population of the study comprised of the students of 9<sup>th</sup> and 10<sup>th</sup> classes studying in rural areas of Lahore. There was a total of 175 public secondary schools in the Lahore district. A total of 24687 students were enrolled in 175 public secondary schools. The list of all secondary schools was obtained from the office of the District Education Officer (DEO). This obtained list served as a sampling frame and through a simple random sampling technique, 20 schools were chosen from a total of 175 schools. The sample size for the study was 160 respondents determined through the online sample size calculator [www.surveysystem.com](http://www.surveysystem.com). Through stratified disproportionate sampling 80 from class 9<sup>th</sup> and 80 from class, 10<sup>th</sup> was chosen thereby making a sample size of 160 respondents. Data were collected through structured, validated and reliable questionnaires whereas the collected data were analyzed using Statistical Package for Social Sciences (SPSS). Descriptive statistics were applied to the data to explore the distribution of data.

## RESULTS

### Demographic Profile of Participants

The table shows that 26.9% of respondents were aged under 15 years, 32.5% were aged 16 and 30.6% of respondents were aged 17 years. Of the total participants of the study, 58.13% were male and 41.87% were females. The respondents were matriculated students and their medium of instruction was regarded as important. Around 69% of respondents were studying in the English medium whereas 31.5% of respondents were going through the Urdu medium. As for as family income was concerned, 18.1% of respondents fell into the income group earning less than 20,000 Rupees in a month. Out of total respondents, 37.5% had earned between 20,000-30,000 Rupees in a month. Around 19% of respondents had earned between 30,000-40,000 and only 15% of respondents reported their family income exceeding 50,000 Rupees in a month.

For the 5.6% of respondents, their guardians didn't have formal education. Of the total respondents, 16.3, 34 and 6.9% of respondents' guardians had primary, matriculation and intermediate level of education, respectively. Among total participants, 36.3% reported their guardian having graduated or above the level of

education. Taking the employment status of guardians, 16.9% were farmers, 43.1% were government employees, 23.8% were working as a labour and 16.3% of respondents had their own business. Regarding the family type, 12.5% were living in nuclear families, 53.8% had a joint family system and 33.8% of respondents had an extended family system.

**Table 1**  
*demographic profile of participants.*

Demographic attributes	N (%)
<b>Age</b>	
<15	43 (26.9)
16	52 (32.5)
17	49 (30.6)
18	12 (7.5)
>19	4 (2.5)
<b>Gender</b>	
Male	93 (58.13)
Female	67 (41.87)
<b>Medium of instruction</b>	
Urdu	50 (31.5)
English	110 (68.7)
<b>Monthly Income</b>	
< 20000	29 (18.1)
20000-30000	60 (37.5)
30000-40000	47 (29.4)
>50000	24 (15.0)
<b>Education of guardian</b>	
Illiterate	9 (5.6)
Primary	26 (16.3)
Matric	56 (35)
Intermediate	11 (6.9)
Graduation or above	58 (36.3)
<b>Employment status</b>	
Farmer	27 (16.9)
Govt. Employee	69 (43.1)
Labor	38 (23.8)
Business	26 (16.3)
<b>Family type</b>	
Nuclear	20 (12.5)
Joint	86 (53.8)
Extended	54 (33.8)

**Knowledge, Attitude and Preventive Practice toward DF**

Table 2 indicates that a large number of respondents (139: 86.9%) had heard about dengue fever. More than half (59.4%) and 90.6% of participants were informed that Anopheles and Aedes were the carrier of dengue facer, respectively. An overwhelming majority of respondents agreed that dengue was different from other mosquitos. This implies that the participants had a good level of awareness regarding the identification of dengue. Around 71% of respondents were informed that dengue is transmittable while 29.4% believed that dengue was not transmittable. This may be related to the sceptical behaviour among participants as for them there was no difference in the common mosquito and dengue.

In the context of awareness regarding breeding sites of dengue, 32 and 56.3% of respondents were informed that clean water and unclean water were the breeding

sites, respectively. Of the total participants, 92.5% were well informed that dengue breeds are in the trash, garbage and used tires. More than one-fifth (22.5%) of respondents were not informed about the breeding sites of the dengue. Regarding common symptoms of dengue fever, information was explored about the high-grade fever, nausea and vomiting, body aches, red spots on the body, pain abdomen, and diarrhea. The majority of respondents (80.6%) were informed about the high-grade fever followed by 73.13% of respondents who were aware of nausea and vomiting. Body aches, red spots on the body, pain abdomen and diarrhea were known in 66.25, 58.12, 38.1 and 20% of participants, respectively. All the symptoms were known to only 16.3% of the total respondents.

Considering the biting time, more than half (51.9%) believed dengue bites in the early morning whereas 75.6% believed its biting time was in the evening. Of the total respondents, 83.1 and 86.1% were informed that dengue bites in the night and noon respectively. Only 3.1% of respondents were unknown of the biting time of dengue.

There were different preventive measures known to study participants. An overwhelming majority (98.8%), 95% and 88.1% of respondents were well informed about the use of mosquito repellent coils, nets and lotions respectively. Only 6.9% of respondents were aware of the mosquito sprays. The awareness regarding wearing protective clothes (98.1%), clean surroundings (95.6%) and use of fire smoke (81.9%) were adequate among the respondents. One in ten respondents (10%) reported doing nothing against the dengue.

**Table 2**  
*Knowledge, attitude and preventive practice (KAP) towards DF.*

	n (%)
<b>Heard about the DF</b>	
Yes	139 (86.9)
No	21 (13.1)
<b>Carrier of dengue</b>	
Anopheles	95 (59.4)
Aedes	145 (90.6)
Different from other mosquitos	145 (90.6)
<b>Dengue is transmittable</b>	
Yes	113 (70.6)
No	47 (29.4)
<b>Breeding site of Dengue</b>	
Clean water	52 (32)
Unclean water	90 (56.3)
Trash/garbage/used tires etc.	148 (92.5)
Don't know	36 (22.5)
<b>A common symptom of dengue fever</b>	
High-grade Fever	129 (80.6)
Nausea and vomiting	117 (73.13)
Body aches	106 (66.25)
Red spots on the body	93 (58.12)
Pain abdomen	61 (38.1)
Diarrhea	32 (20)
All of above	26 (16.3)
<b>Biting time of dengue</b>	

Early morning	83 (51.9)
Evening	121 (75.6)
Night	133 (83.1)
Noon	154 (96.2)
Do not know	5 (3.1)
<b>Preventive measures</b>	
Mosquito repellent coils	158 (98.8)
Mosquito repellent net	152 (95)
Mosquito-repellent lotion	141 (88.1)
Use of mosquito sprays	12 (6.9)
Wear protective clothes	157 (98.1)
Cleaning the surrounding environment	153 (95.6)
Use of fire smoke	115 (81.9)
Do nothing	16 (10)

**Figure 1**  
steps taken after observing the dengue fever.




Figure 1 portrays that 98.8% of respondents were informed to consult the doctor at the earliest the symptoms of dengue are observed. Of the total respondents, 76.3% were informed to visit the nearest Basic Health Unit (BHU) for urgent treatment and prescription. Almost half of the respondents (49.4%) reported self-medication followed by 35.8% of respondents calling to Government Helpline for further guidance and preventive management. Slightly higher than one-fifth (21.2%) of respondents reported their awareness to consult a Hakeem for the treatment.

**Table 3**  
Source of information

Sources	N (%)
School	129 (80.6)
T. V	78 (48.75)
Friends and Family	104 (65)
Doctors	35 (21.87)
Internet	82 (51.25)
Radio	21 (13.12)
Newspapers	84 (52.5)
Flexes and Banners	89 (55.62)
Dengue Aagahi walks	81 (50.6)
Health Teams	48 (30)

**Figure 2**




Table 4 indicates that different information sources played a role in creating awareness among the participants. Around 81% of respondents reported that school contributed to creating awareness. Among all respondents, 48.75, 65, 21.9, 51.3 and 13.1% of respondents reported TV, friends and family, doctors, the internet and radio played their role in creating public awareness. As for as printed media was concerned, 52.5% and 55.62% were informed through newspapers and flexes and banners, respectively. Almost half of the respondents (50.6%) referred their awareness to dengue *aagahi* walks and 30% reported their awareness due to health teams and their awareness campaigns.

**DISCUSSION**

Background variables such as age, education, gender, income, and residential status are regarded as important especially while exploring the knowledge, attitude and skills of the individuals. Similarly, the socio-economic variables were equally important among the population in the context of the incidence of dengue and the execution of preventive measures (Udayanga et al., 2018). Nyamwembe et al. (2018) have reported a positive association between awareness and information utilization. Education is a prominent attribute which can help the individuals in effective utilization of information. In this study, all the respondents were educated and studying 9<sup>th</sup> and 10<sup>th</sup>. This implies that the students had more awareness and understanding of the dengue incidence and preventive measures. Respondents' guardian was also educated and this would have helped the students to become more aware and implement the preventive measures on the household level. Inadequate awareness and lack of knowledge about preventive measures were the key problems that individuals faced (Ho et al., 2013). There was a significant relationship between age, gender and class awareness and knowledge (Adolu et al., 2014). In this study, male and female students participated which reflects the socialization patterns in the area as endorsed by Raudsepp (2001) and Diamontopoulos et al. (2003). At the household level, females often take all the managerial, sweeping and cleaning activities, thus they had a very prominent role in dengue prevention. These findings are supported by those of Ero et al. (2001) as

they have reported an association between demographic attributes like age, gender, and education to perform different maintenance tasks at home. Adeolu et al. (2014) also found a relationship between demographic attributes i.e. gender, age, class and their level of awareness and practices of waste management. This is an endorsement of current findings that demographic attributes have key significance in waste management which could be the breeding site for dengue.

In this study majority of the respondents were informed about dengue, its breeding and its transmission though many of the respondents didn't have a basic awareness about dengue, its transmission, breeding and the carrier of dengue fever. With special reference to the breeding of the dengue, respondents had more inclination towards breeding in dirty and unclean water. In a true sense, dengue breeds in fresh water and half of the respondents were unknown of this fact. These results are similar to those of Hossain et al. (2021) as they found that respondents had misconceptions regarding the breeding of dengue in dirty and sewage water. They perceived that swage drains and unclean water were the common breeding sites of the dengue. Our study confirmed that there were different symptoms of dengue fever and some of the respondents were well informed about the identification of symptoms. Although, fever was the foremost and obvious reason for the majority of the respondents. It was perceived that a large number of respondents who participated in this study had not personally experienced dengue fever not any of their family members embraced the dengue, therefore the

respondents didn't have full confidence in typical symptoms of the dengue.

In our study, we further explored the respondents who were informed to use mosquito repellent oil, net, lotion, wear protective clothing and clean the environment to prevent dengue. Findings are almost consistent with those of Hossain et al. (2021) as they found that most of the respondents were using mosquito coils to prevent dengue. However, our findings are not comparable with those of Acharya et al. (2005) who affirmed that cleaning the surroundings and environment did not help prevent dengue. Considering the results of our study, we believe that cleaning the surrounding can help in destroying the breeding sites of the mosquitos. Once the mosquito sites are destroyed the mosquito would not be able to breed and as result, the population of dengue will remain in control. Our study also confirmed the use of indigenous techniques like using smoke to prevent dengue.

## CONCLUSION

Dengue has become a key threat and dengue fever is the mosquito born disease that is lethal in its impact and its prevention and remedy are inevitable. Being aware about the dengue fever and its prevention seem important especially in the process of dengue infestation. Although, the awareness and knowledge among respondents was adequate enough but still the prevention strategies were not fully implemented and adopted. Negligible role of modern media gadgets in creating dengue awareness was the significant gap observed in this study.

## REFERENCES

- Adeolu, A. T., Enesi, D. O., & Adeolu, M. O. (2014). Assessment of secondary school students' knowledge, attitude and practice towards waste management in Ibadan, Oyo State, Nigeria. *Journal of Research in Environmental Science and Toxicology*, 3(5), 66-73.
- Ali, H., Alvi, A., Fatima, S., Zafar, F., Naveed, S., Khan, K., ... & Mallick, N. (2017). Dengue fever in Pakistan, episodes of epidemic to endemic: treatment challenges, prevention and current facts. *J Bioequiv Availab*, 9(5), 473-476. <https://doi.org/10.4172/jbb.1000347>
- Arora, P., Arora, M., Sharma, V., & Kotwal, A. (2017). Dengue: Awareness, preventive practices and water storage behaviour in an urban community of Delhi. *International Journal Of Community Medicine And Public Health*, 4(12), 4460. <https://doi.org/10.18203/2394-6040.ijcmph20175148>
- Brady, O. J., Gething, P. W., Bhatt, S., Messina, J. P., Brownstein, J. S., Hoen, A. G., Moyes, C. L., Farlow, A. W., Scott, T. W., & Hay, S. I. (2012). Refining the global spatial limits of dengue virus transmission by evidence-based consensus. *PLoS Neglected Tropical Diseases*, 6(8), e1760. <https://doi.org/10.1371/journal.pntd.0001760>
- Degallier, N. P. T. V., Vilarinhos, P. D. R., de Carvalho, M. L., Knox, M. B., & Caetano Jr, J. (2000). People's knowledge and practice about dengue, its vectors, and control means in Brasilia (DF), Brazil: its relevance with entomological factors. *Journal of the American Mosquito Control Association*, 16(2), 114-123. [https://www.biodiversitylibrary.org/content/part/JAMCA/JAMCA\\_V16\\_N2\\_P114-123.pdf](https://www.biodiversitylibrary.org/content/part/JAMCA/JAMCA_V16_N2_P114-123.pdf)
- Dhar-Chowdhury, P., Emdad Haque, C., Michelle Driedger, S., & Hossain, S. (2014). Community perspectives on dengue transmission in the city

- of Dhaka, Bangladesh. *International Health*, 6(4), 306-316. <https://doi.org/10.1093/inthealth/ihu032>
- Dhimal, M., Aryal, K. K., Dhimal, M. L., Gautam, I., Singh, S. P., Bhusal, C. L., & Kuch, U. (2014). Knowledge, attitude and practice regarding dengue fever among the healthy population of Highland and lowland communities in central Nepal. *PLoS ONE*, 9(7), e102028. <https://doi.org/10.1371/journal.pone.0102028>
- Diamantopoulos, A., Schlegelmilch, B. B., Sinkovics, R. R., & Bohlen, G. M. (2003). Can socio-demographics still play a role in profiling green consumers? A review of the evidence and an empirical investigation. *Journal of Business Research*, 56(6), 465-480. [https://doi.org/10.1016/s0148-2963\(01\)00241-7](https://doi.org/10.1016/s0148-2963(01)00241-7)
- Harapan, H., Rajamoorthy, Y., Anwar, S., Bustamam, A., Radiansyah, A., Angraini, P., Fasli, R., Salwiyadi, S., Bastian, R. A., Oktiviyari, A., Akmal, I., Iqbalamin, M., Adil, J., Henrizal, F., Darmayanti, D., Pratama, R., Setiawan, A. M., Mudatsir, M., Hadisoemarto, P. F., ... Müller, R. (2018). Knowledge, attitude, and practice regarding dengue virus infection among inhabitants of Aceh, Indonesia: A cross-sectional study. *BMC Infectious Diseases*, 18(1). <https://doi.org/10.1186/s12879-018-3006-z>
- Ho, T., Huang, M., Wang, S., Hsu, H., & Liu, C. (2013). Knowledge, attitude, and practice of dengue disease among healthcare professionals in southern Taiwan. *Journal of the Formosan Medical Association*, 112(1), 18-23. <https://doi.org/10.1016/j.jfma.2012.11.004>
- Hossain, M. I., Alam, N. E., Akter, S., Suriea, U., Aktar, S., Shifat, S. K., Islam, M. M., Aziz, I., Islam, M. M., Islam, M. S., & Mohiuddin, A. K. (2021). Knowledge, awareness and preventive practices of dengue outbreak in Bangladesh: A countrywide study. *PLOS ONE*, 16(6), e0252852. <https://doi.org/10.1371/journal.pone.0252852>
- Islam, Z., Mohanan, P., Bilal, W., Hashmi, T., Rahmat, Z., Abdi, I., Riaz, M. M., & Essar, M. Y. (2022). Dengue virus cases surge amidst COVID-19 in Pakistan: Challenges, efforts and recommendations. *Infection and Drug Resistance*, 15, 367-371. <https://doi.org/10.2147/idr.s347571>
- Jahan, F. (2011). Dengue Fever (DF) in Pakistan. *Asia Pacific Family Medicine*, 10(1), 1. <https://doi.org/10.1186/1447-056x-10-1>
- Jeelani, S., Sabesan, S., & Subramanian, S. (2015). Community knowledge, awareness and preventive practices regarding dengue fever in Puducherry – South India. *Public Health*, 129(6), 790-796. <https://doi.org/10.1016/j.puhe.2015.02.026>
- Lambrechts, L., Scott, T. W., & Gubler, D. J. (2010). Consequences of the expanding global distribution of aedes albopictus for dengue virus transmission. *PLoS Neglected Tropical Diseases*, 4(5), e646. <https://doi.org/10.1371/journal.pntd.0000646>
- Mayxay, M., Cui, W., Thammavong, S., Khensakhou, K., Vongxay, V., Inthasoum, L., Sychareun, V., & Armstrong, G. (2013). Dengue in Peri-urban pak-ngum district, Vientiane capital of Laos: A community survey on knowledge, attitudes and practices. *BMC Public Health*, 13(1). <https://doi.org/10.1186/1471-2458-13-434>
- Naing, C., Ren, W. Y., Man, C. Y., Fern, K. P., Qiqi, C., Ning, C. N., & Ee, C. W. (2011). Awareness of dengue and practice of dengue control among the semi-urban community: A cross sectional survey. *Journal of Community Health*, 36(6), 1044-1049. <https://doi.org/10.1007/s10900-011-9407-1>
- Nyamwembe, E. O., Wamutitu, J., & Tanui, E. (2018). Relationship Between Students' Perceptions And Utilization Of Open Educational Resources For Academic Work In Private Universities In Kenya. *International Journal of Education and Research*, 6(9), 113-128.
- Olli, E., Grendstad, G., & Wollebaek, D. (2001). Correlates of environmental behaviors. *Environment and Behavior*, 33(2), 181-208. <https://doi.org/10.1177/0013916501332002>
- Sharmin, S., Viennet, E., Glass, K., & Harley, D. (2015). The emergence of dengue in Bangladesh: Epidemiology, challenges and future disease risk. *Transactions of The Royal Society of Tropical Medicine and Hygiene*, 109(10), 619-627. <https://doi.org/10.1093/trstmh/trv067>
- Siddiqua, M., Alam, A. N., Muraduzzaman, A., & Shirin, T. (2018). NS-1 antigen positive dengue infection and molecular characterization of dengue viruses in a private medical college Hospitalin Dhaka, Bangladesh. *Bangladesh Journal of Medical Science*, 17(4), 669-673. <https://doi.org/10.3329/bjms.v17i4.38334>

- Udayanga, L., Gunathilaka, N., Iqbal, M. C., Pahalagedara, K., Amarasinghe, U. S., & Abeyewickreme, W. (2018). Socio-economic, knowledge attitude practices (KAP), household related and demographic based appearance of non-dengue infected individuals in high dengue risk areas of kandy district, Sri Lanka. *BMC Infectious Diseases*, 18(1). <https://doi.org/10.1186/s12879-018-2995-y>
- Udayanga, L., Gunathilaka, N., Iqbal, M. C., Pahalagedara, K., Amarasinghe, U. S., & Abeyewickreme, W. (2018). Socio-economic, knowledge attitude practices (KAP), household related and demographic based appearance of non-dengue infected individuals in high dengue risk areas of kandy district, Sri Lanka. *BMC Infectious Diseases*, 18(1). <https://doi.org/10.1186/s12879-018-2995-y>
- Vachvanichsanong, P., Thisyakorn, U., & Thisyakorn, C. (2015). Dengue hemorrhagic fever and the kidney. *Archives of Virology*, 161(4), 771-778. <https://doi.org/10.1007/s00705-015-2727-1>
- Waheed, S. (2021). *Dengue fever*. World Health Organization - Regional Office for the Eastern Mediterranean. <http://www.emro.who.int/pak/programmes/dengue-fever.html>
- WHO. (2021, December 14). *Dengue fever – Pakistan*. [Www.who.int. https://www.who.int/emergencies/disease-outbreak-news/item/dengue-fever-pakistan](https://www.who.int/emergencies/disease-outbreak-news/item/dengue-fever-pakistan)